

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/51131948>

2-(4-Methylanilino)acetohydrazide

Article in *Acta Crystallographica Section E Structure Reports Online* · September 2009

DOI: 10.1107/S1600536809033169 · Source: PubMed

CITATIONS

0

READS

15

5 authors, including:



Shridhar Ashok Malladi

Gogte Institute of Technology

111 PUBLICATIONS 398 CITATIONS

[SEE PROFILE](#)



Mahesh Padaki

Jain University

125 PUBLICATIONS 1,179 CITATIONS

[SEE PROFILE](#)



Arun M Islor

National Institute of Technology Karnataka

547 PUBLICATIONS 4,009 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Medicinal organic chemistry [View project](#)



Hydrogels for tissue engineering [View project](#)

2-(4-Methylanilino)acetohydrazide

**Hoong-Kun Fun,^{a,*‡} Chin Sing Yeap,^{a,§} Shridhar Malladi,
Mahesh Padaki^b and Arun M. Isloor^b**

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Chemistry, National Institute of Technology–Karnataka, Surathkal, Mangalore 575 025, India
Correspondence e-mail: hkfun@usm.my


Received 17 August 2009; accepted 20 August 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.039; wR factor = 0.110; data-to-parameter ratio = 15.9.

In the title molecule, $\text{C}_9\text{H}_{13}\text{N}_3\text{O}$, the non-hydrogen atoms of the hydrazide group are essentially planar [maximum deviation = 0.028 (1) \AA for one of the N atoms]. The mean plane of this group forms a dihedral angle of 83.34 (5) $^\circ$ with the plane of the benzene ring. In the crystal structure, molecules are linked by intermolecular $\text{N}-\text{H}\cdots\text{O}$, $\text{N}-\text{H}\cdots\text{N}$ and weak $\text{C}-\text{H}\cdots\text{N}$ hydrogen bonds into a two-dimensional network parallel to the ab plane. Additional stabilization is provided by a weak $\text{C}-\text{H}\cdots\pi$ interaction.

Related literature

For the biological activity of hydrazide derivatives, see: Ozdemir *et al.* (2009); Khattab (2005). For synthetic applications, see: Isloor *et al.* (2009); Holla & Udupa (1992). For a related structure, see: Zhang & Shi (2009). For the stability of the temperature controller used for the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

$\text{C}_9\text{H}_{13}\text{N}_3\text{O}$	$c = 15.4756 (4)\text{ \AA}$
$M_r = 179.22$	$\alpha = 87.002 (2)^\circ$
Triclinic, $P\bar{1}$	$\beta = 84.282 (2)^\circ$
$a = 5.1481 (1)\text{ \AA}$	$\gamma = 82.849 (2)^\circ$
$b = 5.9262 (2)\text{ \AA}$	$V = 465.76 (2)\text{ \AA}^3$

‡ Thomson Reuters ResearcherID: A-3561-2009.
§ Thomson Reuters ResearcherID: A-5523-2009.

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.09\text{ mm}^{-1}$

$T = 100\text{ K}$
 $0.52 \times 0.15 \times 0.07\text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.956$, $T_{\max} = 0.994$

11697 measured reflections
2703 independent reflections
2301 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.110$
 $S = 1.03$
2703 reflections
170 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.34\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.20\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N1 \cdots O1 ⁱ	0.875 (17)	2.162 (17)	3.0271 (11)	170.0 (13)
N2—H1N2 \cdots N3 ⁱⁱ	0.887 (17)	2.287 (16)	3.0302 (12)	141.3 (13)
N3—H1N3 \cdots O1 ⁱⁱⁱ	0.901 (14)	2.252 (13)	3.0614 (11)	149.4 (13)
N3—H2N3 \cdots O1 ^{iv}	0.914 (14)	2.281 (15)	3.0889 (12)	147.1 (12)
C7—H7B \cdots N3 ^v	0.991 (15)	2.545 (14)	3.4341 (14)	149.2 (11)
C9—H9B \cdots Cg ^v	0.96 (2)	2.94 (2)	3.7469 (16)	142.3 (14)

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x, -y + 1, -z + 1$; (iii) $-x + 1, -y, -z + 1$; (iv) $x - 1, y, z$; (v) $-x + 1, -y + 1, -z + 1$. Cg is the centroid of the C1–C6 benzene ring.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINT* (Bruker, 2005); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

HKF thanks Universiti Sains Malaysia (USM) for the Research University Golden Goose grant No. 1001/PFIZIK/811012. CSY thanks USM for the award of a USM Fellowship. AMI is grateful to the Head of the Department of Chemistry and Director, NITK, Surathkal, India, for providing research facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2884).

References

- Bruker (2005). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- Holla, B. S. & Udupa, K. V. (1992). *Farmaco*, **47**, 305–318.
- Isloor, A. M., Kalluraya, B. & Shetty, P. (2009). *Eur. J. Med. Chem.* **44**, 3784–3787.
- Khattab, S. N. (2005). *Molecules*, **10**, 1218–1228.
- Ozdemir, A., Turan-Zitouni, G., Kaplançikli, Z. A. & Tunali, Y. (2009). *J. Enzyme Inhibit. Med. Chem.* **24**, 825–831.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Zhang, Y.-X. & Shi, Z.-Q. (2009). *Acta Cryst. E* **65**, o1538.

supplementary materials

Acta Cryst. (2009). E65, o2234 [doi:10.1107/S1600536809033169]

2-(4-Methylanilino)acetohydrazide

H.-K. Fun, C. S. Yeap, S. Malladi, M. Padaki and A. M. Isloor

Comment

In organic chemistry, hydrazides are a class of organic compounds sharing a common functional group characterized by a nitrogen to nitrogen covalent bond with 4 substituents with at least one of them being an acyl group. They are also starting materials for many heterocycles including 1,2,4-triazoles (Isloor *et al.*, 2009; Holla & Udupa, 1992). Many hydrazide derivatives have showed significant biological activities (Ozdemir *et al.*, 2009; Khattab, 2005). In view of the biological and synthetic importance of hydrazides, we hereby report the crystal structure of the title compound (I).

The bond lengths and angles of the title compound (I), (Fig. 1) are comparable to its related structure (Zhang & Shi, 2009). A maximum deviation of 0.028 (1) Å for atom N2 from the mean plane of the hydrazide group form by atoms O1, N2, N3, C7 and C8 indicates that it is essentially coplanar. The mean plane of the hydrazide makes dihedral angle of 83.34 (5)° with C1–C6 benzene ring. In the crystal structure, the molecules are linked by intermolecular N1—H1N1···O1ⁱ, N2—H1N2···N3ⁱⁱ, N3—H1N3···O1ⁱⁱⁱ, N3—H2N3···O1^{iv} and C7—H7B···N3^v (see Table 1 for symmetry codes) hydrogen bonds into two-dimensional network parallel to *ab* plane (Fig. 2, Table 1). The crystal structure is also stabilized by a C—H···π interaction (Table 1).

Experimental

Ethyl [(4-methylphenyl)amino]acetate (19.3 g, 0.1 mol) and hydrazine hydrate (99%, 0.2 mol) in ethanol (200 ml) was heated on a water-bath for 6 h. Excess of ethanol was removed by distillation. On cooling, colourless block-shaped single crystals of 2-[(4-methylphenyl)amino]acetohydrazide begin to separate (Holla & Udupa, 1992). It was collected by filtration and recrystallized from ethanol. Yield: 13.2 g, 73.7%, *M.p.* 423–426 K.

Refinement

All hydrogen atoms were located from the difference Fourier map and refined freely, with N—H = 0.876 (15)–0.912 (15) Å; C—H = 0.96 (2)–1.04 (2) Å.

Figures




Fig. 1. The molecular structure of the title compound with atom labels and 50% probability ellipsoids for non-H atoms.

supplementary materials




Fig. 2. Part of the crystal structure of (I), viewed along the b axis, showing the two-dimensional network parallel to ab plane. Intermolecular hydrogen bonds are shown as dashed lines.

2-(4-Methylanilino)acetohydrazide

Crystal data

$C_9H_{13}N_3O$	$Z = 2$
$M_r = 179.22$	$F_{000} = 192$
Triclinic, $P\bar{1}$	$D_x = 1.278 \text{ Mg m}^{-3}$
Hall symbol: -P 1	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 5.1481 (1) \text{ \AA}$	Cell parameters from 6230 reflections
$b = 5.9262 (2) \text{ \AA}$	$\theta = 2.7\text{--}31.2^\circ$
$c = 15.4756 (4) \text{ \AA}$	$\mu = 0.09 \text{ mm}^{-1}$
$\alpha = 87.002 (2)^\circ$	$T = 100 \text{ K}$
$\beta = 84.282 (2)^\circ$	Block, colourless
$\gamma = 82.849 (2)^\circ$	$0.52 \times 0.15 \times 0.07 \text{ mm}$
$V = 465.76 (2) \text{ \AA}^3$	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	2703 independent reflections
Radiation source: fine-focus sealed tube	2301 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.026$
$T = 100 \text{ K}$	$\theta_{\max} = 30.0^\circ$
φ and ω scans	$\theta_{\min} = 2.7^\circ$
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	$h = -7 \rightarrow 7$
$T_{\min} = 0.956$, $T_{\max} = 0.994$	$k = -8 \rightarrow 8$
11697 measured reflections	$l = -21 \rightarrow 21$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.039$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.110$	$w = 1/[\sigma^2(F_o^2) + (0.0559P)^2 + 0.1344P]$
$S = 1.03$	where $P = (F_o^2 + 2F_c^2)/3$
2703 reflections	$(\Delta/\sigma)_{\max} < 0.001$
	$\Delta\rho_{\max} = 0.34 \text{ e \AA}^{-3}$

170 parameters

 $\Delta\rho_{\min} = -0.20 \text{ e } \text{\AA}^{-3}$ Primary atom site location: structure-invariant direct
methods

Extinction correction: none

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.63005 (14)	0.15409 (11)	0.39934 (5)	0.01915 (17)
N1	0.43188 (17)	0.74133 (14)	0.33320 (6)	0.01783 (18)
N2	0.26039 (16)	0.39523 (13)	0.43209 (5)	0.01650 (18)
N3	0.12716 (17)	0.23345 (14)	0.48445 (6)	0.01765 (18)
C1	0.1124 (2)	0.95874 (17)	0.24809 (7)	0.0231 (2)
C2	-0.0573 (2)	0.9751 (2)	0.18340 (8)	0.0282 (2)
C3	-0.0748 (2)	0.7920 (2)	0.13190 (7)	0.0280 (2)
C4	0.0857 (2)	0.5915 (2)	0.14806 (7)	0.0257 (2)
C5	0.2577 (2)	0.57064 (17)	0.21309 (7)	0.0208 (2)
C6	0.27314 (19)	0.75502 (16)	0.26432 (6)	0.01770 (19)
C7	0.62171 (19)	0.54434 (16)	0.34553 (7)	0.01795 (19)
C8	0.50373 (18)	0.34635 (15)	0.39460 (6)	0.01517 (18)
C9	-0.2624 (3)	0.8132 (3)	0.06212 (9)	0.0405 (3)
H1A	0.116 (3)	1.091 (3)	0.2848 (10)	0.031 (4)*
H2A	-0.167 (3)	1.117 (3)	0.1751 (11)	0.042 (4)*
H4A	0.081 (3)	0.457 (3)	0.1131 (10)	0.032 (4)*
H5A	0.363 (3)	0.424 (3)	0.22235 (10)	0.031 (4)*
H7A	0.713 (3)	0.484 (2)	0.2916 (9)	0.024 (3)*
H7B	0.757 (3)	0.592 (2)	0.3797 (9)	0.026 (3)*
H9A	-0.199 (4)	0.895 (4)	0.0098 (14)	0.070 (6)*
H9B	-0.435 (4)	0.878 (3)	0.0840 (13)	0.064 (6)*
H9C	-0.290 (4)	0.655 (4)	0.0400 (14)	0.072 (6)*
H1N1	0.485 (3)	0.869 (3)	0.3461 (9)	0.029 (4)*
H1N2	0.185 (3)	0.538 (3)	0.4343 (9)	0.028 (3)*
H1N3	0.250 (3)	0.125 (2)	0.5030 (9)	0.023 (3)*
H2N3	0.031 (3)	0.167 (2)	0.4485 (9)	0.028 (4)*

supplementary materials

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0173 (3)	0.0139 (3)	0.0257 (4)	0.0010 (2)	-0.0024 (3)	-0.0012 (3)
N1	0.0208 (4)	0.0112 (3)	0.0219 (4)	-0.0032 (3)	-0.0028 (3)	-0.0005 (3)
N2	0.0155 (4)	0.0110 (3)	0.0223 (4)	-0.0013 (3)	0.0001 (3)	0.0014 (3)
N3	0.0159 (4)	0.0135 (4)	0.0233 (4)	-0.0029 (3)	-0.0015 (3)	0.0035 (3)
C1	0.0247 (5)	0.0183 (5)	0.0246 (5)	0.0002 (4)	0.0004 (4)	0.0021 (4)
C2	0.0246 (5)	0.0294 (6)	0.0277 (6)	0.0031 (4)	-0.0008 (4)	0.0081 (4)
C3	0.0224 (5)	0.0410 (6)	0.0209 (5)	-0.0083 (4)	-0.0023 (4)	0.0072 (4)
C4	0.0274 (5)	0.0303 (5)	0.0210 (5)	-0.0109 (4)	-0.0013 (4)	-0.0011 (4)
C5	0.0225 (5)	0.0183 (4)	0.0216 (5)	-0.0041 (3)	-0.0006 (4)	-0.0007 (3)
C6	0.0178 (4)	0.0160 (4)	0.0189 (4)	-0.0033 (3)	0.0006 (3)	0.0015 (3)
C7	0.0156 (4)	0.0158 (4)	0.0222 (5)	-0.0027 (3)	-0.0004 (3)	0.0010 (3)
C8	0.0156 (4)	0.0140 (4)	0.0165 (4)	-0.0020 (3)	-0.0032 (3)	-0.0019 (3)
C9	0.0291 (7)	0.0679 (10)	0.0258 (6)	-0.0130 (6)	-0.0079 (5)	0.0127 (6)

Geometric parameters (\AA , $^\circ$)

O1—C8	1.2421 (11)	C2—H2A	0.962 (17)
N1—C6	1.3995 (13)	C3—C4	1.3867 (17)
N1—C7	1.4437 (12)	C3—C9	1.5094 (17)
N1—H1N1	0.876 (15)	C4—C5	1.3961 (15)
N2—C8	1.3313 (12)	C4—H4A	0.988 (15)
N2—N3	1.4205 (11)	C5—C6	1.3978 (14)
N2—H1N2	0.886 (15)	C5—H5A	0.979 (15)
N3—H1N3	0.902 (14)	C7—C8	1.5213 (13)
N3—H2N3	0.912 (15)	C7—H7A	0.978 (14)
C1—C2	1.3850 (16)	C7—H7B	0.989 (14)
C1—C6	1.4022 (13)	C9—H9A	0.97 (2)
C1—H1A	0.993 (15)	C9—H9B	0.96 (2)
C2—C3	1.3963 (18)	C9—H9C	1.04 (2)
C6—N1—C7	120.37 (8)	C4—C5—C6	120.25 (10)
C6—N1—H1N1	115.9 (10)	C4—C5—H5A	119.3 (9)
C7—N1—H1N1	113.7 (10)	C6—C5—H5A	120.4 (9)
C8—N2—N3	122.60 (8)	C5—C6—N1	122.90 (9)
C8—N2—H1N2	120.9 (10)	C5—C6—C1	118.06 (10)
N3—N2—H1N2	115.5 (10)	N1—C6—C1	118.97 (9)
N2—N3—H1N3	107.5 (9)	N1—C7—C8	113.36 (8)
N2—N3—H2N3	106.4 (9)	N1—C7—H7A	114.3 (8)
H1N3—N3—H2N3	107.4 (12)	C8—C7—H7A	106.5 (8)
C2—C1—C6	120.65 (10)	N1—C7—H7B	107.1 (8)
C2—C1—H1A	119.7 (9)	C8—C7—H7B	108.4 (8)
C6—C1—H1A	119.6 (9)	H7A—C7—H7B	107.0 (12)
C1—C2—C3	121.82 (10)	O1—C8—N2	123.22 (9)
C1—C2—H2A	118.2 (10)	O1—C8—C7	121.37 (9)
C3—C2—H2A	120.0 (10)	N2—C8—C7	115.40 (8)

C4—C3—C2	117.19 (10)	C3—C9—H9A	113.1 (13)
C4—C3—C9	121.88 (12)	C3—C9—H9B	111.3 (12)
C2—C3—C9	120.92 (12)	H9A—C9—H9B	111.5 (17)
C3—C4—C5	122.03 (10)	C3—C9—H9C	112.3 (12)
C3—C4—H4A	120.2 (9)	H9A—C9—H9C	103.7 (17)
C5—C4—H4A	117.8 (9)	H9B—C9—H9C	104.4 (17)
C6—C1—C2—C3	−0.26 (17)	C7—N1—C6—C1	−172.79 (9)
C1—C2—C3—C4	−0.06 (17)	C2—C1—C6—C5	0.35 (15)
C1—C2—C3—C9	179.61 (11)	C2—C1—C6—N1	−176.67 (9)
C2—C3—C4—C5	0.28 (16)	C6—N1—C7—C8	−83.08 (11)
C9—C3—C4—C5	−179.38 (11)	N3—N2—C8—O1	2.92 (14)
C3—C4—C5—C6	−0.19 (16)	N3—N2—C8—C7	−176.34 (8)
C4—C5—C6—N1	176.77 (9)	N1—C7—C8—O1	169.09 (8)
C4—C5—C6—C1	−0.13 (15)	N1—C7—C8—N2	−11.63 (12)
C7—N1—C6—C5	10.34 (14)		


Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1N1···O1 ⁱ	0.875 (17)	2.162 (17)	3.0271 (11)	170.0 (13)
N2—H1N2···N3 ⁱⁱ	0.887 (17)	2.287 (16)	3.0302 (12)	141.3 (13)
N3—H1N3···O1 ⁱⁱⁱ	0.901 (14)	2.252 (13)	3.0614 (11)	149.4 (13)
N3—H2N3···O1 ^{iv}	0.914 (14)	2.281 (15)	3.0889 (12)	147.1 (12)
C7—H7B···N3 ^v	0.991 (15)	2.545 (14)	3.4341 (14)	149.2 (11)
C9—H9B···Cg ^{iv}	0.96 (2)	2.94 (2)	3.7469 (16)	142.3 (14)

Symmetry codes: (i) $x, y+1, z$; (ii) $-x, -y+1, -z+1$; (iii) $-x+1, -y, -z+1$; (iv) $x-1, y, z$; (v) $-x+1, -y+1, -z+1$.

supplementary materials

Fig. 1



supplementary materials

Fig. 2

